

ACTUAL PROBLEMS OF DISEASES OF THE ENDOCRINE SYSTEM

Mullacheri Kamalon Rohith Krishnan
Fergana Medical Institute of Public Health

Annotation: The endocrine system is made up of glands that make hormones. Hormones are the body's chemical messengers. They carry information and instructions from one set of cells to another. The endocrine system coordinates the functions of various organs through hormones, which are chemicals that are released into the bloodstream from specialized cells found inside the endocrine glands (endocrine glands). Endocrine glands release hormones into the bloodstream. This lets the hormones travel to cells in other parts of the body. The endocrine hormones help control mood, growth and development, the way our organs work, metabolism, and reproduction[1,2].

Key words: Growth and development, hypothalamus, pituitary, thyroid, parathyroids, adrenals, pineal body, endocrine system.

INTRODUCTION: The endocrine system regulates how much of each hormone is released. This can depend on levels of hormones already in the blood, or on levels of other substances in the blood, like calcium. Many things affect hormone levels, such as stress, infection, and changes in the balance of fluid and minerals in blood[3].

While many parts of the body make hormones, the major glands that make up the endocrine system are the: hypothalamus, pituitary, thyroid, parathyroids, adrenals, pineal body, the ovaries, the testes.

The pancreas is part of the endocrine system and the digestive system. That's because it secretes hormones into the bloodstream, and makes and secretes enzymes into the digestive tract. While in the blood, hormones influence the function of target tissues, which can be either another endocrine gland or a receptor.

Some hormones act on the cells of the same organ that is their source (paracrine effect), and sometimes even on the cells themselves that produce them (autocrine effect)[5,6]. Endocrine system disorders result from disruption of the endocrine glands and/or their target tissues.

MATERIALS AND METHODS OF RESEARCH: Analysis of statistical indicators of morbidity and mortality from diseases of the endocrine system obtained from WHO reporting statistics from 2021. Too much or too little of any hormone can harm the body. Medicines can treat many of these problems.

The endocrine system is a messenger system in an organism comprising feedback loops of hormones that are released by internal glands directly into the circulatory system and that target and regulate distant organs. In vertebrates, the hypothalamus is the neural control center for all endocrine systems[7]. In humans, the major endocrine glands are the thyroid, parathyroid, pituitary, pineal, and adrenal glands, and the (male) testicles and (female) ovaries. The hypothalamus, pancreas, and thymus also function as endocrine glands, among other functions. (The hypothalamus and pituitary glands are organs of the neuroendocrine system. One of the most important functions of the hypothalamus—it is located in the brain adjacent to the pituitary gland—is to link the endocrine system to the nervous system via the pituitary gland.) Other organs, such as the kidneys, also have roles within the endocrine system by secreting certain hormones. The study of the endocrine system and its disorders is known as endocrinology[8].

There are many endocrine glands in your body that release different hormones. Many of these glands are controlled by the hypothalamus and the pituitary gland (also known as the master gland) in our brain. There are many medical conditions that can affect your glands and cause hormonal

problems. The proportion of patients with pathology of the endocrine system is constantly increasing in all economically developed countries of the world[9,10]. Diabetes mellitus occupies a leading place in the structure of all endocrine diseases, which today is one of the most dangerous challenges to the global community and an important priority for national healthcare systems. The second place in the structure of endocrine pathology belongs to various diseases of the thyroid gland. One of the leading indicators that determine the epidemiological situation of diabetes mellitus in the world is the prevalence due to the growth in the size and age of the population, urbanization of the territory, the increase in the prevalence of obesity and a sedentary lifestyle."

According to experts, in the European region, which includes 59 countries, the number of people with diabetes is 59.3 million, or 9.5% of the adult population. The highest rate is observed in Turkey - 15.8%. The prevalence of diabetes mellitus in Germany is 12.95%, in Spain - 10.53%, Italy - 7.55%, France - 7.20%, in the UK - 6.97%. The minimum national prevalence of diabetes mellitus is observed in Azerbaijan - 2.78%, China has the largest number of people with diabetes mellitus - 109.6 million, and the prevalence rate reaches 11.6%. The majority of the region's population lives in North America and the Caribbean in the USA, Mexico and Canada, and in these same countries the vast majority of people with diabetes are registered. The national prevalence of diabetes mellitus in Mexico is 11.7%, Canada - 11.21%, USA - 10.90%. Another important indicator that determines the epidemiological situation with regard to diabetes mellitus is the incidence[11,12]. There are clear geographic differences in growth trends, but the average annual growth rate is 3%. It has been established that type 1 diabetes in the world affects 81,100 children under the age of 15 every year. Of the 499,100 children living with type 1 diabetes today, 27% are in Europe, where the latest and most reliable data on the incidence of new diabetes cases is available, and 23% are in North America and the Caribbean[13]. Global health care spent about \$549 billion in 2017 to prevent complications of diabetes and treat the disease itself. This figure is expected to exceed US\$628 billion by 2035.

Diseases of the endocrine system are common, including conditions such as diabetes mellitus, thyroid disease, and obesity. Endocrine disease is characterized by misregulated hormone release (a productive pituitary adenoma), inappropriate response to signaling (hypothyroidism), lack of a gland (diabetes mellitus type 1, diminished erythropoiesis in chronic kidney failure), or structural enlargement in a critical site such as the thyroid (toxic multinodular goitre). Hypofunction of endocrine glands can occur as a result of loss of reserve, hyposecretion, agenesis, atrophy, or active destruction[14,15]. Hyperfunction can occur as a result of hypersecretion, loss of suppression, hyperplastic or neoplastic change, or hyperstimulation.

Endocrinopathies are classified as primary, secondary, or tertiary. Primary endocrine disease inhibits the action of downstream glands. Secondary endocrine disease is indicative of a problem with the pituitary gland. Tertiary endocrine disease is associated with dysfunction of the hypothalamus and its releasing hormones.

As the thyroid, and hormones have been implicated in signaling distant tissues to proliferate, for example, the estrogen receptor has been shown to be involved in certain breast cancers. Endocrine, paracrine, and autocrine signaling have all been implicated in proliferation, one of the required steps of oncogenesis[16].

Other common diseases that result from endocrine dysfunction include Addison's disease, Cushing's disease and Graves' disease. Cushing's disease and Addison's disease are pathologies involving the dysfunction of the adrenal gland. Dysfunction in the adrenal gland could be due to primary or secondary factors and can result in hypercortisolism or hypocortisolism[17]. Cushing's disease is characterized by the hypersecretion of the adrenocorticotrophic hormone (ACTH) due to a pituitary

adenoma that ultimately causes endogenous hypercortisolism by stimulating the adrenal glands. Some clinical signs of Cushing's disease include obesity, moon face, and hirsutism. Addison's disease is an endocrine disease that results from hypocortisolism caused by adrenal gland insufficiency. Adrenal insufficiency is significant because it is correlated with decreased ability to maintain blood pressure and blood sugar, a defect that can prove to be fatal.

Graves' disease involves the hyperactivity of the thyroid gland which produces the T3 and T4 hormones. Graves' disease effects range from excess sweating, fatigue, heat intolerance and high blood pressure to swelling of the eyes that causes redness, puffiness and in rare cases reduced or double vision. The amount of spending on diabetes treatment varies significantly from region to region and from country to country. Only 1/5 of global spending occurs in middle- and low-income countries, although about 80% of patients with diabetes live in them. An average of 356 US dollars per year is spent per patient in middle- and low-income countries, and 5,621 US dollars in high-income countries. Thyroid pathology ranks second in prevalence after diabetes among all endocrinopathies. Since the beginning of the 90- In the 1990s, it is also characterized by an increase in prevalence, depending on many factors, such as gender, age, genetic predisposition, the presence of goitrogens in food (thioglycosides, thiocyanates), and iodine supply of the region.

The number of patients who have suffered temporary and permanent disability due to thyroid pathology is increasing. The incidence of thyroid cancer is currently increasing throughout the world. The rate of increase in the detection of this disease is 4% per year. Today, this pathology is the most common malignant neoplasm of the endocrine system, accounting for 2.2% in the structure of cancer incidence. A common condition is hypothyroidism. Manifest hypothyroidism occurs in 0.2-2% of the population, subclinical - 4-10% (in the elderly - 7-26%). The prevalence of hypothyroidism ranges from 3 to 16% in men and from 4 to 21% in women and increases with age. Such a high prevalence determines the medical and social significance of hypothyroidism. The most common cause of this pathology is autoimmune thyroiditis. In general, autoimmune thyroid diseases occur in 2-5% of the population.

Thyrotoxicosis remains a serious problem, leading to a worsening life prognosis and accompanied by an increase in mortality from diseases of the cardiovascular system. Thus, mortality from coronary heart disease, heart failure, arrhythmias, valve defects and arterial hypertension in combination with thyrotoxicosis increased 1.2 times compared to the general population. The reason for this is the development of changes in the cardiovascular system. Among the manifestations of "thyrotoxic heart" are pulmonary hypertension, diastolic dysfunction, the development of heart failure, dilatation of the heart cavities, atrial fibrillation and left ventricular myocardial hypertrophy. Atrial fibrillation, which occurs in 2-25% of cases with thyrotoxicosis, often persists even after thyrotoxicosis has been eliminated.

One of the most common diseases of the thyroid gland is diffuse toxic goiter, the prevalence of which in the population is 1-3%, the incidence is from 5 to 23 cases per 100 thousand population per year, the ratio of men to women is 1:5 - 1:7. Diffuse toxic goiter is primarily dangerous due to disorders of the cardiovascular system associated with thyrotoxicosis and leading to a worsening prognosis of the disease and loss of ability to work.

CONCLUSIONS: The data presented indicate the high significance of endocrine pathology - diabetes mellitus and thyroid diseases. Thus, the nature of the prevalence and course of the disease, the frequency of identified patients depend on the level and quality of local endocrinological care, timely active identification of patients and complications, provision of sufficient quantities of drugs, proper organization of dispensary observation and therapeutic measures.

REFERENCES:

1. Anatomy of the Endocrine System. www.hopkinsmedicine.org. November 19, 2019. Retrieved June 14, 2022.
2. Gardner, Shoback (2017). Greenspan's Basic and Clinical Endocrinology (10th ed.). McGraw Hill / Medical. pp. 49–68. ISBN 978-1259589287.
3. Jump up to: Marieb E (2014). Anatomy & physiology. Glenview, IL: Pearson Education, Inc. ISBN 978-0-321-86158-0.
4. Sherwood, L. (1997). Human Physiology: From Cells to Systems. Wadsworth Pub Co. ISBN 978-0-495-39184-5.
5. Гинзбург М. М. Ожирение: влияние на развитие метаболического синдрома. Профилактика и лечение/ Гинзбург М. М., Н.И. Крюков – М. : Медпрактика, 2002.
6. IN Raxmatjonovna. Qabziyat sabablari, tashxislash va davolash. научные исследования и общественные проблемы 1 (1), 205-207
7. НР Исакова, Ж Юнусов, АГ Худоярова. Возможные пути коррекции секреторной функции поджелудочной железы с её протоковой системы. неделя науки-2017, 489-491
8. Isaqova, N.R. Influence of constipation on anthropometric indicators of children. Science and Innovation, Volume 1, Issue 8, pp. 888-892, 2022.
9. IN Raxmatjonovna. The most pressing problem today is iodine deficiency. World Bulletin of Public Health 23, 97-100
10. Isakova N.R. The effect of constipation due to diseases of the colon on the anthropometric parameters of children. Asian journal of multidimensional research, Volume:10, Issue 5, pp. 666-669
11. IN Raxmatjonovna. Effects of colonic diseases on children's health. World bulletin of public health 23, 101-103, 2023
12. ИН Рахматжоновна. Влияние запора на антропометрические показатели детей при заболеваниях толстого кишечника. Тиббиётда янги кун 2 (34), 85-87
13. Y.Nishonov., A.Abdulhakimov., N.Madrahimova. Scientific bases of methods for studying anthropometry of the eye bowl. Science and Innovation, Volume 1, Issue 8, pp. 1001-1006, 2022.
14. 7-18 ёшли болаларнинг кўз косаси антропометриясини ўрганиш. Ю.Н.Нишонов., А.Р.Абдулхакимов., Н.Р.Мадрахимова. Scientific impulse 1(5), 910-913, 2022.
15. Palvanova M.S. Morphological changes in the bone tissue of the child's body in the age aspect. World Bulletin of Public Health, 94-96, 2023
16. Р.Т.Юсупова, О.Е. Шаланкова Репродуктивное здоровье девочек-подростков, проживающих в условиях Ферганской долины. Университетская наука: взгляд в будущее, 612-614, 2020
17. Palvanova M.S., Akhmatov B.K. Chronic myeloid leukemia epidemiology in the Fergana region over decade from 2010 until 2020. Science and innovation, Volume1, issue 8, pp. 1020-1025