

Volume10, Issue08, Aug-2023.

SJIF 2019: 4.702 2020: 4.737 2021: 5.071 2022: 4.919 2023: 6.980

---

## ANATOMICAL AND ETIOLOGICAL FEATURES OF SPINAL CORD TUMORS THAT ARE NOW COMMON

**Djalalov Ravshanbek**

Dosent of Namangan State University medical faculty , Uzbekistan

**Annotation :** A spinal tumor is a growth that develops within your spinal canal or within the bones of your spine. A spinal cord tumor, also called an intradural tumor, is a spinal tumor that begins within the spinal cord or the covering of the spinal cord (dura). A tumor that affects the bones of the spine (vertebrae) is called a vertebral tumor. Spinal cord tumors may be classified as one of three different types depending on where they occur relative to the protective membranes of the spinal cord.

**Key words :** Spinal , Signs , tumor , shok , growth .

---

Spinal cord tumors can cause different signs and symptoms, especially as tumors grow. The tumors may affect your spinal cord or the nerve roots, blood vessels or bones of your spine. Signs and symptoms may include:

- Pain at the site of the tumor due to tumor growth
- Back pain, often radiating to other parts of your body
- Feeling less sensitive to pain, heat and cold
- Loss of bowel or bladder function
- Difficulty walking, sometimes leading to falls
- Back pain that's worse at night
- Loss of sensation or muscle weakness, especially in your arms or legs
- Muscle weakness , which may be mild or severe, in different parts of your body

Back pain is a common early symptom of spinal tumors. Pain may also spread beyond your back to your hips, legs, feet or arms and may worsen over time — even with treatment.

Spinal tumors progress at different rates depending on the type of tumor

Volume10, Issue08, Aug-2023.

SJIF 2019: 4.702 2020: 4.737 2021: 5.071 2022: 4.919 2023: 6.980


There are many causes of back pain, and most back pain isn't caused by a tumor. But because early diagnosis and treatment are important for spinal tumors, see your doctor about your back pain if:

- It's persistent and progressive
- It's not activity related
- It gets worse at night
- You have a history of cancer and develop new back pain
- You have other symptoms of cancer, such as nausea, vomiting or dizziness

Seek immediate medical attention if you experience:

- Progressive muscle weakness or numbness in your legs or arms
- Changes in bowel or bladder function

Causes



© MAYO FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH. ALL RIGHTS RESERVED.

Sp

inal anatomyOpen pop-up dialog box

Volume10, Issue08, Aug-2023.

SJIF 2019: 4.702 2020: 4.737 2021: 5.071 2022: 4.919 2023: 6.980

---

It's not clear why most spinal tumors develop. Experts suspect that defective genes play a role. But it's usually not known whether such genetic defects are inherited or simply develop over time. They might be caused by something in the environment, such as exposure to certain chemicals. In some cases, however, spinal cord tumors are linked to known inherited syndromes, such as neurofibromatosis 2 and von Hippel-Lindau disease.

#### Risk factors


Spinal cord tumors are more common in people who have:

- **Neurofibromatosis 2.** In this hereditary disorder, benign tumors develop on or near the nerves related to hearing. This may lead to progressive hearing loss in one or both ears. Some people with neurofibromatosis 2 also develop spinal canal tumors.
- **Von Hippel-Lindau disease.** This rare, multisystem disorder is associated with blood vessel tumors (hemangioblastomas) in the brain, retina and spinal cord and with other types of tumors in the kidneys or adrenal glands.

#### Complications

Spinal tumors can compress spinal nerves, leading to a loss of movement or sensation below the location of the tumor. This can sometimes cause changes in bowel and bladder function. Nerve damage may be permanent.

However, if caught early and treated aggressively, it may be possible to prevent further loss of function and regain nerve function. Depending on its location, a tumor that presses against the spinal cord itself may be life-threatening.



#### Nervous system [Open pop-up dialog box](#)

It's not clear why most spinal tumors develop. Experts suspect that defective genes play a role. But it's usually not known whether such genetic defects are inherited or simply develop over

Volume10, Issue08, Aug-2023.

SJIF 2019: 4.702 2020: 4.737 2021: 5.071 2022: 4.919 2023: 6.980

---

time. They might be caused by something in the environment, such as exposure to certain chemicals. In some cases, however, spinal cord tumors are linked to known inherited syndromes, such as neurofibromatosis 2 and von Hippel-Lindau disease.

Risk factors

Spinal cord tumors are more common in people who have:

- **Neurofibromatosis 2.** In this hereditary disorder, benign tumors develop on or near the nerves related to hearing. This may lead to progressive hearing loss in one or both ears. Some people with neurofibromatosis 2 also develop spinal canal tumors.
- **Von Hippel-Lindau disease.** This rare, multisystem disorder is associated with blood vessel tumors (hemangioblastomas) in the brain, retina and spinal cord and with other types of tumors in the kidneys or adrenal glands.

**References :**

1. Andreev A. Yu. Mikhailova L. M. Starnov A. A. Closure of the Ca<sup>2+</sup> - dependent CsA pore: the role of Md<sup>2+</sup>, adenine nucleotides, and the conformational state of ADP/ATR-antiporter. *Biochemistry*. 1994. T. 59 c1589-1597.
2. Podymova S. D. Liver diseases. Moscow: Meditsina, 1984, 480S.
3. Leniger A. Bioenergetics and metabolism. *Fundamentals of biochemistry*. Moscow: Nauka, 1985. T-2. pp. 439-550.
4. Mayer K. P. Hepatitis and the consequences of hepatitis. Moscow. 2001. p. 423.
5. Panda S.V., Blanchaer M.C. reversible inhibition of mitochondrial adenosine diphosphate phosphorylation by longchain acyl coenzyme A esters // *J. Biol. Chem.* 1971. Vol. 246. N-2. P. 402-411.
6. Tareev E. M. Tareev I. E. Chronic hepatitis and cirrhosis. // In the book: "Manual of internal diseases, ed. Tareeva E. M. Moscow, 1965, vol. 5.
7. Khanin M. N. Experimental heliotropic hepatitis. // The second All-Union Conference of Pathophysiologists. Theses of reports, Kiev, 1963, p. 25