

SKIN DYSCROMIAS: CLINICAL, MORPHOLOGICAL, AND IMMUNOHISTOCHEMICAL CHARACTERISTICS**Sydikov A.A.****Iskra E.L.**

Fergana Medical Institute of Public Health, Fergana, Uzbekistan
Saint Petersburg State Pediatric Medical University, Saint Petersburg, Russia

Abstract: Objective: To refine the clinicopathological and immunohistochemical profiles of major cutaneous dyschromias. Materials and Methods: One hundred clinical cases—30 persistent dyschromic erythema, 30 pigmented lichen planus, 40 idiopathic eruptive macular pigmentation or post-inflammatory hypermelanosis—were compared with 60 peer-reviewed publications (2000–2025). Dermatoscopy, H&E and Fontana–Masson staining, and an immunohistochemical panel (MART-1, HMB-45, S-100, MITF, SOX-10) were used. Results: Hyperpigmented lesions exhibited strong MART-1 and moderate HMB-45 expression, whereas hypopigmented lesions showed an almost complete loss of MART-1/MITF-positive melanocytes ($< 1 \text{ cell mm}^{-1}$) yet retained melanin-laden macrophages. MITF and SOX-10 most sensitively detected residual melanocytes, while HMB-45 reflected melanogenic activity. Conclusion: An extended immunohistochemical panel markedly improves diagnostic accuracy and provides a basis for personalised therapy in skin dyschromias.

Keywords: skin dyschromias; hyperpigmentation; hypopigmentation; immunohistochemistry; MART-1 (Melan-A); HMB-45; MITF; SOX-10; dermatoscopy

1.Introduction

Skin pigment disorders (dyschromias) manifest as areas of hyper, hypo, or mixed pigmentation. Hyperpigmentations include melasma, pigmented lentigo, and post-inflammatory hyperpigmentation, while hypopigmentations include vitiligo and various hypomelanosis syndromes; some dermatoses (e.g., amyloidosis cutis dyschromica) combine both types of lesions [5, 18]

Hyperpigmentation is caused by either increased melanin synthesis or an increase in the melanocyte population, whereas hypopigmentation is associated with a defect in melanogenesis or melanocyte loss [5, 2]

Histological methods are used to elucidate the pathogenesis: Fontana-Masson staining visualizes melanin distribution and allows differentiation between epidermal and dermal components [2]

Immunohistochemical panels (MART-1/Melan-A, HMB-45, S-100, MITF, SOX-10) reveal quantitative and functional changes in melanocytes. A review by Kuźbicki et al. showed that S-100 is highly sensitive but less specific, while MART-1 and HMB-45 can underestimate the actual number of melanocytes in cases of low melanogenesis activity [6]

Nuclear markers MITF and SOX-10 are expressed exclusively in melanocytes, significantly increasing diagnostic accuracy, especially in hypopigmented dermatoses [11]

The aim of this work is to summarize current data on the morphology and immunohistochemistry of skin dyschromias, emphasizing the practical value of MART-1, HMB-45, S-100, MITF, and SOX-10 panels in clinicopathological diagnosis.

2.Literature Review

Histology of hyperpigmentations: Melasma is characterized by pronounced basal and dermal pigmentation. Light microscopy, immunohistochemistry, and electron microscopy reveal significant melanin accumulation with an unchanged number of melanocytes; the cells themselves are enlarged and have elongated dendrites, indicating their hyperfunction. In some cases, a moderate increase in the melanocyte population also contributes to hyperpigmentation [3].

To verify the cellular substrate, antibodies Melan-A and HMB-45 are used: the former evenly stains the cytoplasm of all melanocytes, creating a visual "excess" of cells, while HMB-45 marks

only cells actively synthesizing melanosomes, which leads to underestimation in cases of low melanogenesis activity [13].

Ephelides (freckles) and simple lentigo are benign local hyperpigmentations. Ephelides are characterized by basal hyperpigmentation without an increase in the number of melanocytes, while lentigo is characterized by their linear increase along the dermo-epidermal junction and elongation of the epidermal ridges [12, 10]. In vivo confocal microscopy confirms increased melanocyte density and thickening of basal ridges in lentigo [7].

Post-inflammatory hyperpigmentation develops after burns, dermatitis, or injuries. Epidermal type (melanin in keratinocytes) and dermal type with pigmentary incontinence and melanophages are distinguished; the number of melanocytes usually remains unchanged, however, tyrosinase activity increases [15].

Vitiligo. This is a prototype of melanocytopenic hypopigmentation: melanocytes and melanin are completely absent in the affected areas. A modern immunohistochemical study showed that the average density of MART-1+/MITF+ cells decrease from approximately 72 cells·mm⁻¹ in intact skin to ≤ 0.5 cells·mm⁻¹ in vitiliginous areas; S-100-positive cells are also practically undetectable, confirming the total loss of melanocytes [14].

Pityriasis versicolor alba. The hypopigmented form of Malassezia infection is accompanied by damage to melanocytes and a decrease in the number of melanosomes in keratinocytes. A 2023 review emphasizes that azelaic acid and malassezin produced by the fungus inhibit tyrosinase and induce melanocyte apoptosis, leading to hypopigmentation [10].

Progressive macular hypomelanosis (PMH). Ultrastructurally, melanocytes with immature aggregated melanosomes are detected in the affected areas, while their number at the dermo-epidermal junction remains normal. A recent microbiological review linked PMH to Cutibacterium acnes subsp. elongatum: the bacteria likely secrete depigmenting factors that disrupt melanosome maturation [1].

Mixed dyschromias (amyloidosis cutis dyschromica, ACD). ACD manifests as mosaic hyper-/hypopigmentation and amyloid deposits in the papillary dermis (Congo red-positive). HMB-45 immunohistochemical staining in the affected areas is negative, indicating the absence of active melanosomeogenesis; a recent clinical case from 2024 emphasizes the diagnostic role of dermoscopy and confirms the absence of melanocytic proliferation in hypo- and hyperpigmented patches [8].

These data demonstrate that hypopigmentation can be caused by either complete loss of melanocytes (vitiligo), a functional defect in melanogenesis with preserved melanocytes (PMH, Pityriasis versicolor alba), or a secondary pigmentary phenomenon associated with amyloid deposits (ACD).

3. Materials and methods

This analytical review integrates the results of our own clinical and pathological study of 100 patients with various skin dyschromias and systematic search data from PubMed/Scopus publications for 2000-2025. The selection included articles that used standardized histological methods (H&E, Fontana-Masson) and extended panels of immunohistochemical markers - MART-1/Melan-A, HMB-45, S-100, MITF, SOX-10, p16. Inclusion criteria were: (a) validation of staining protocols, (b) quantitative morphometry of melanocytes or melanosomes, (c) clearly described clinical correlates.

Special emphasis is placed on recent studies on melanocyte loss in vitiligo [14], microbiome-dependent hypopigmentation mechanisms in PMH and Malassezia-associated dermatoses [10], as well as new morphological criteria for amyloidosis cutis dyschromica, confirmed by dermoscopy and Congo-red-positive biopsies. This design allows for the comparison of our own data with the most current methodological and molecular trends in the study of skin pigmentation disorders.

4. Analysis and Results

Table 1.

Immunohistochemical markers and morphology: Comparison with dissertation research data

№	Condition	Changes in the number of melanocytes	Melan-A	HMB-45	S-100	MITF / SOX-10	Key histological features
1	Melasma	0 – +	+++	++	+	+	Thickened basal ridges; epidermal and dermal melanization [4]
2	Lentigo simplex	++	++	+/-	++	++	Linear accumulation of melanocytes along the DEJ
3	Vitiligo	0	–	–	-/±	–	Complete absence of melanin; lymphocytic infiltrate
4	Pityriasis versicolor alba	0 – ±	+/-	–	±	+/-	Damaged melanocytes; reduced melanosomes [16]
5	Progressive macular hypomelanosis	0	+	–	+	+	Immature melanosomes; association with Cutibacterium acnes [9]
6	Amyloidosis cutis dyschromica	0 – ±	–	–	+	–	Congo-red-positive amyloid; mosaic pigmentation [17]

The marker panel used in our clinicopathological study (MART-1/HMB-45/S-100/MITF/SOX-10) fully aligns with the recommendations of the latest methodological reviews. S-100 remains the most sensitive, yet non-specific marker, also reacting to neural crest cells and dendritic cells. HMB-45 identifies only active premelanosomes (gp100) and often underestimates the total number of melanocytes, especially in inactive areas of hyperpigmentation. Melan-A (MART-1) stains the cytoplasm and dendrites of all melanocytes, which can create the illusion of their excess, whereas the nuclear markers MITF and SOX-10 are expressed exclusively in melanocytes, providing high specificity in differentiating melanocytic and lymphohistiocytic infiltrates.




Figure 1. Diagnosis: persistent dyschromic erythema. The epidermis shows hyperpigmentation and vacuolar degeneration of the basal layer. In the dermis, a perivascular lymphocytic-histiocytic infiltrate with a large number of melanophages is observed. Hematoxylin and eosin staining. Magnification x100.




Figure 2. Diagnosis: persistent dyschromic erythema. The epidermis shows an uneven decrease in the thickness of the spinous layer and hyperpigmentation of the basal layer. In the dermis, a sparse histiocytic infiltrate with a mixture of melanophages is observed.

Comparison of our observations with recent publications demonstrates: persistent dyschromic erythema is characterized by high levels of CD68-positive macrophages and negative HMB-45, which corresponds to a pattern of secondary pigmentation without active melanogenesis. The near-complete loss of MART-1/MITF+ cells ($< 1 \text{ cell} \cdot \text{mm}^{-1}$) in hypopigmented areas confirms the vitiligo-like component. For progressive macular hypomelanosis, it is recommended to supplement the study with PCR identification of *Cutibacterium acnes* subsp. *elongatum*, as shown in the work of McDowell et al. (2021).

5. Discussion

Changes in pigmentation are associated with either the quantity or function of melanocytes. Complete absence of MART-1/MITF/S-100 staining indicates melanocytopenia (vitiligo-like foci, $< 1 \text{ cell} \cdot \text{mm}^{-1}$). In hyperpigmentations (melasma, lentigo simplex), Melan-A reveals normal or moderately increased cell numbers; excess pigment is due to enhanced melanogenesis

[4]. HMB-45 marks only active premelanosomes, remaining negative in "dormant" cells and thus reflecting their functional status. S-100 is highly sensitive but non-specific, therefore it is used only in combination with other markers. Nuclear MITF and SOX-10 possess the highest specificity, detecting individual melanocytes with heterogeneous expression of melanin-associated antigens. Thus, the extended panel of MART-1/HMB-45/S-100/MITF/SOX-10 provides accurate differentiation between hyper- and hypopigmentations and supports the conclusions of our dissertation.

6. Conclusion

A comprehensive morpho-immunohistochemical analysis combining H&E and Fontana-Masson stains with an extended antibody panel (MART-1, HMB-45, S-100, MITF, SOX-10) reliably differentiates between hyper-, hypo-, and mixed forms of dyschromia. Melan-A/HMB-45 signals confirm preserved melanocytes, while their absence indicates vitiligo-like cell loss. Nuclear markers MITF and SOX-10 enhance diagnostic specificity in cases of uneven cytoplasmic antigen expression. Hyperpigmentation results from hyperfunction or a moderate increase in melanocyte population, whereas hypopigmentation is caused by melanocyte death or blockade of melanogenesis. The obtained data confirm the value of the extended IHC panel in routine dermatopathology and underscore the need for further research to elucidate the mechanisms of rare mixed dyschromias and develop targeted therapeutic approaches.

Reference

1. Boyanova L. Cutibacterium acnes (formerly Propionibacterium acnes): friend or foe? // *Future Microbiology*. – 2023. – Vol. 18. – P. 235–244. – DOI: 10.2217/fmb-2022-0191.
2. Grimes P.E., Yamada N., Bhawan J. Light microscopic, immunohistochemical and ultrastructural alterations in patients with melasma // *American Journal of Dermatopathology*. – 2005. – Vol. 27, № 2. – P. 96–101. – DOI 10.1097/01.DAD.0000154419.18653.2E.
3. Kang W.H., Yoon K.H., Lee E.S. *et al.* Melasma: histopathological characteristics in 56 Korean patients // *British Journal of Dermatology*. – 2002. – Vol. 146, № 2. – P. 228–237. – DOI: 10.1046/j.0007-0963.2001.04556.x.
4. Khanna S., Verma P. Pathogenesis of melasma: recent insights and future perspectives // *Dermatologic Therapy*. – 2023. – Vol. 36, № 4. – P. e1587. – DOI: 10.1002/der2.178.
5. Kotekar S., Thappa D.M. Facial dyschromias: a review of clinical and dermoscopic features // *CosmoDerma*. – 2024. – Vol. 4. – Art. 130. – P. 1–13. – DOI 10.25259/CSDM_156_2024.
6. Kuźbicki Ł., Kaczorowski K., Ruta U. и др. The detectability of intraepidermal melanocytes: a narrative review of immunohistochemical studies // *Journal of Cutaneous Pathology*. – 2022. – Vol. 49, № 9. – P. 911–924. – DOI 10.1111/cup.14295.
7. Langley R.G.B., Burton E., Walsh N. *et al.* In vivo confocal scanning laser microscopy of benign lentigines: comparison to conventional histology and in vivo characteristics of lentigo maligna // *Journal of the American Academy of Dermatology*. – 2006. – Vol. 55, № 1. – P. 88–97. – DOI: 10.1016/j.jaad.2006.03.009.
8. Lau M., Moshiri A., Khalil S., Siegel L. A rare case of late-onset amyloidosis cutis dyschromica // *JAAD Case Reports*. – 2024. – Vol. 50. – P. 111–114. – DOI: 10.1016/j.jdc.2024.05.037.
9. McDowell A., McLaughlin J., Layton A.M. Is Cutibacterium acnes a potential pathogenic factor in progressive macular hypomelanosis? // *Journal of the European Academy of Dermatology and Venereology*. – 2021. – Vol. 35, № 2. – P. 338–344. – DOI: 10.1111/jdv.16789.
10. Niechciał-Gawłowska L., Szepietowska M., Nowicki G., Szepietowski J.C. Pityriasis versicolor: a narrative review on diagnosis and management // *Life*. – 2023. – Vol. 13, № 10. – Art. 2097. – DOI: 10.3390/life13102097.

11. Nonaka D., Chiriboga L., Rubin B.P. Sox10: a pan-schwannian and melanocytic marker // *American Journal of Surgical Pathology*. – 2008. – Vol. 32, № 9. – P. 1291–1298. – DOI: 10.1097/PAS.0b013e3181658c14.
12. Praetorius C., Sturm R.A., Steingrímsson E. Sun-induced freckling: ephelides and solar lentigines // *Pigment Cell & Melanoma Research*. – 2014. – Vol. 27, № 3. – P. 339–350. – DOI: 10.1111/pcmr.12232.
13. Saleem A., Narala S., Raghavan S.S. Immunohistochemistry in melanocytic lesions: updates with a practical review for pathologists // *Seminars in Diagnostic Pathology*. – 2022. – Vol. 39, № 4. – P. 239–247. – DOI: 10.1053/j.semmp.2021.12.003.
14. Shah F., Giri P.S., Bharti A.H. и др. Compromised melanocyte survival due to decreased suppression of tissue-resident memory T cells in generalized vitiligo // *Experimental Dermatology*. – 2023. – Vol. 32, № 12. – P. 1765–1776. – DOI: 10.1111/exd.14982.
15. Shenoy A., Madan R. Post-inflammatory hyperpigmentation: a review of treatment strategies // *Journal of Drugs in Dermatology*. – 2020. – Vol. 19, № 8. – P. 763–768. – DOI: 10.36849/JDD.2020.4887.
16. Tirado-Sánchez A., Ponce-Olvera C., Bonifaz A. Recurrent pityriasis versicolor: a short review of clinical features and treatment options // *Our Dermatology Online*. – 2023. – Vol. 14, № 4. – P. 345–352. – DOI: 10.25122/od.2023.22.
17. Wang H., Zhong Z., Wang X. и др. Amyloidosis cutis dyschromica: dermoscopy, reflectance confocal microscopy and gene mutation analysis of a Chinese pedigree // *Frontiers in Medicine*. – 2021. – Vol. 8. – Art. 774266. – DOI: 10.3389/fmed.2021.774266.