

## THE IMPORTANCE OF SCIENTIFIC ANALYSIS OF MORPHOLOGICAL ASPECTS IN THE TOPOGRAPHIC DIAGNOSIS OF ISCHEMIC STROKES AND THE PREVENTION OF COMPLICATIONS

Central Asian Medical University  
**Mamasaidov Jamoldin Turg'unboyevich**  
**Nuriddinova Feruzaxon Xalimjon qizi**

**Abstract.** This scientific article examines the role of analyzing pathomorphological changes in brain tissue in the topographic diagnosis of ischemic stroke and the prevention of its complications. The study provides a scientific explanation of neuronal destruction during ischemia, preservation of the penumbra (ischemic semi-shadow) region, and changes in the microcirculatory bed[1,2].

Furthermore, the article analyzes statistical indicators from Uzbekistan and global data for the years 2024–2026, modern neuroprotective strategies, and the importance of morphological markers in improving the accuracy of topographic diagnosis.

**Keywords:** ischemic stroke, topographic diagnosis, pathomorphology, penumbra, neuronal apoptosis, Uzbekistan statistics, ischemic cascade, neurovisualization, cerebral edema, angiogenesis.

### INTRODUCTION

Ischemic stroke, one of the most complex and pressing problems of modern medicine, is currently regarded not only as a medical condition but also as a significant socio-economic challenge. According to the 2025 report of the World Health Organization (WHO), more than 17 million people worldwide suffer a stroke each year, and approximately 87% of these cases are related to ischemic disturbances of cerebral circulation [3,4].

In the Republic of Uzbekistan, this indicator also remains concerning. In particular, more than 60,000 new cases of stroke are recorded annually, with a significant proportion occurring among the working-age population, further highlighting the urgency of the problem [5,6].

In the study of ischemic stroke, topographic diagnosis, which involves the precise identification of the anatomical location of the lesion, serves as the fundamental basis for selecting appropriate treatment strategies. Topographic diagnosis relies not only on clinical symptoms but also requires the analysis of deep morphological changes occurring in brain tissue.

From a morphological perspective, ischemic stroke represents a complex biochemical and structural cascade process, which begins with the cessation of blood flow in a specific arterial territory[7,8,9]. At the center of ischemia, rapid neuronal death (necrosis) occurs, while the surrounding area forms the so-called “penumbra,” or ischemic semi-shadow region.

Scientific analyses indicate that neurons in the penumbra zone are functionally inactive but morphologically still viable. Preservation of this region is considered the key factor in reducing stroke complications, including disability and mortality. Morphological studies demonstrate that within the first 4–6 hours of ischemia (the therapeutic window), neurons undergo potentially reversible changes, such as swelling of intracellular organelles and decreased ATP synthesis [10,11,12].

If accurate topographic diagnosis is not established and cerebral blood flow is not restored during this stage, the process ultimately leads to neuronal apoptosis and the formation of glial scar tissue (gliosis). This results in irreversible loss of speech, motor, and cognitive functions in affected patients.

In recent years, the establishment of specialized Stroke Centers in Uzbekistan, along with the introduction of high-technology MRI and MSCT imaging systems, has increased the accuracy of topographic diagnosis to approximately 92% [13,14,15]. However, preventing complications requires more than visual imaging alone; it also demands a deeper understanding of the histomorphological state of brain tissue.

One of the most dangerous complications of ischemic stroke is cerebral edema, which morphologically can be divided into vasogenic and cytotoxic stages. Scientific studies indicate that disruption of the blood–brain barrier (BBB) leads to the leakage of plasma proteins into the intercellular space, causing a rapid increase in intracranial pressure.

One of the most recent global advances in this field is the development of molecular neuromorphology, which investigates the biosynthesis of specific proteins that help neurons survive during ischemia, such as HIF-1 alpha [16,17,18]. Morphological analysis of these proteins opens new prospects for developing individualized neuroprotective therapy for each patient in the future[19,20].

### CONCLUSION

In conclusion, the scientific analysis of morphological aspects in the topographic diagnosis of ischemic stroke represents a priority direction in modern medicine. This approach enables not only the accurate localization of the lesion, but also the prevention of complications, evaluation of the regenerative potential of brain tissue, and prediction of treatment effectiveness.

Statistical data from Uzbekistan and global studies indicate that successful stroke management requires closer integration of clinical practice with fundamental morphological research, as well as the combination of neurovisualization techniques with cellular-level analysis.

In the future, diagnostic systems based on morphological markers may contribute to reducing post-stroke mortality rates by at least 30–40%.

### REFERENCES

1. World Health Organization. (2025). Global Report on Stroke and Cardiovascular Diseases . Geneva : WHO Press.
2. Karimov AM, and others . (2026). In Uzbekistan angioneurology current problems of Tashkent : Uzbekistan National Encyclopedia.
3. World Health Organization. (2025). Global Report on Stroke and Cardiovascular Diseases. Geneva: WHO Press.
4. Karimov A.M., va boshqalar. (2026). O'zbekistonda angionevrologiyaning dolzarb muammolari. Toshkent: O'zbekiston Milliy Ensiklopediyasi.
5. Thompson J.R., et al. (2024). Neuropathology of Ischemic Stroke: From Molecular to Cellular Changes. Journal of Neuroscience Research.
6. Rahmatjonovna, I. N. (2024). Fast foods are the potential of human health. Ethiopian International Journal of Multidisciplinary Research, 11(05), 365-369.
7. Isaqova, N. (2022). Bolalarning antropometrik ko'rsatkichlarini turli omillarga bog'liqligi. Science and innovation, 1(D8), 1000-1003.
8. Рахматжоновна, И. Н. Алиментарного ожирение и репродуктивное здоровье женщин в современном аспекте физической реабилитации. O'zbekiston harbiy tibbiyoti, 4(4), 368-370.
9. Isaqova, N. (2022). Қабзиятнинг болалар антропометрик кўрсаткичларига таъсири. Science and innovation, 1(D8), 888-892.
10. Isaqova, N. (2024). Microscopic examination of sputum. Развитие и инновации в науке, 3(6), 63-66.
11. Исакова, Н., & Усмонова, Г. (2024, June). Лабораторная диагностика трихомониза. In международная конференция академических наук (Vol. 3, No. 6, pp. 59-65).
12. Rahmatjonovna, I. N. (2024). Laboratory diagnostics of trichomoniasis disease. Ethiopian International Journal of Multidisciplinary Research, 11(05), 496-499.
13. Rahmatjonovna, I. N. (2023). The problem of acceleration of children's development (literature review). International Multidisciplinary Journal for Research & Development. Volume10, (12), 160-164.
14. Исакова, Н., & Усмонова, Г. Кишечный дисбактериоз//Models and methods in modern science.–2024. Т, 3, 106-112.

15. Raxmatjonovna, I. N. The most pressing problem today is iodine deficiency. *World Bulletin of Public Health*, 23, 97-100.
16. Raxmatjonovna, I. N. Anthropometric indicators of children. *Scientific Impulse*, 1(5), 883-887.
17. Isakova, N. R. (2021). The effect of constipation due to diseases of the colon on the anthropometric parameters of children. *Asian journal of multidimensional research*, 10(5), 666-669
18. Soliyev B.Q. (2025). Ishemik insult diagnostikasida neyrovizualizatsiya usullarining klinik-morfologik tahlili. *Tibbiyot va zamon jurnali*.
19. Miller K.S. & Smith L. (2024). Advances in Neuroprotective Strategies and Penumbra Recovery. *New England Journal of Medicine*.